

Polymorphic Automorphisms and the Picard Group

Pieter Hofstra, Jason Parker, Philip Scott

Brandon University, Manitoba, Canada

Formal Structures for Computation and Deduction 2021
June 28, 2021

Introduction

- The notion of *definable automorphism* occurs throughout algebra, model theory, and computer science.
- In first-order logic, an automorphism α of a model M of a first-order theory is *definable* if there is a formula $\varphi(x, y)$ such that $\alpha(a) = b$ iff $M \models \varphi(a, b)$ for all $a, b \in M$. E.g. if G is a group, then the inner automorphism induced by $g \in G$ is definable by the formula $y = gxg^{-1}$.
- Definable automorphisms are *polymorphic* or *uniform*, and can provide a generalized notion of *inner automorphism*.

Motivation

- To motivate this, we recall that George Bergman proved in [1] that the definable group automorphisms, i.e. the inner automorphisms given by conjugation, can be characterized purely *categorically* as the automorphisms that extend naturally along any group homomorphism.
- To see this, observe first that if α is an inner automorphism of a group G (induced by $s \in G$), then for each group morphism $f : G \rightarrow H$ with domain G we can ‘push forward’ α to define an inner automorphism

$$\alpha_f : H \xrightarrow{\sim} H$$

by conjugation with $f(s) \in H$ (so that $\alpha_{\text{id}_G} = \alpha$).

Motivation

- This family of automorphisms $(\alpha_f)_f$ is *coherent*, in the sense that it satisfies the following *naturality* property: if $f : G \rightarrow G'$ and $f' : G' \rightarrow G''$ are group homomorphisms, then the following diagram commutes:

$$\begin{array}{ccc} G' & \xrightarrow{\alpha_f} & G' \\ f' \downarrow & & \downarrow f' \\ G'' & \xrightarrow{\alpha_{f' \circ f}} & G'' \end{array}$$

Bergman's Theorem

For a group G , let us call an *arbitrary* family of automorphisms

$$\left(\alpha_f : \mathbf{cod}(f) \xrightarrow{\sim} \mathbf{cod}(f) \right)_{\mathbf{dom}(f)=G}$$

with the above naturality property an *extended inner automorphism* of G . Such a family is a natural automorphism of $G/\mathbf{Group} \rightarrow \mathbf{Group}$.

Theorem (Bergman [1])

Let G be a group and $\alpha : G \xrightarrow{\sim} G$ an automorphism of G . Then α is an **inner** automorphism of G iff there is an extended inner automorphism $(\alpha_f)_f$ of G with $\alpha = \alpha_{\mathbf{id}_G}$.

This provides a completely *categorical* characterization of inner automorphisms of groups: they are exactly those group automorphisms that are 'coherently extendible' along morphisms out of the domain.

Covariant Isotropy

- We have a functor $\mathcal{Z} : \mathbf{Group} \rightarrow \mathbf{Group}$ that sends any group G to its group of extended inner automorphisms $\mathcal{Z}(G)$. We refer to \mathcal{Z} as the *covariant isotropy group (functor)* of the category \mathbf{Group} . (Bergman's theorem actually entails that $\mathcal{Z} \cong \text{Id} : \mathbf{Group} \rightarrow \mathbf{Group}$.)
- In fact, *any* category \mathbb{C} has a *covariant isotropy group (functor)*

$$\mathcal{Z}_{\mathbb{C}} : \mathbb{C} \rightarrow \mathbf{Group}$$

that sends each object $C \in \mathbb{C}$ to the group of extended inner automorphisms of C , i.e. families of automorphisms

$$\left(\alpha_f : \mathbf{cod}(f) \xrightarrow{\sim} \mathbf{cod}(f) \right)_{\mathbf{dom}(f)=C}$$

in \mathbb{C} with the same naturality property as before, i.e. natural automorphisms of the projection functor $C/\mathbb{C} \rightarrow \mathbb{C}$.

Covariant Isotropy

- We can also turn Bergman's characterization of inner automorphisms in **Group** into a *definition* of inner automorphisms in an arbitrary category \mathbb{C} : if $C \in \mathbb{C}$, we say that an automorphism $\alpha : C \xrightarrow{\sim} C$ is *inner* if there is an extended inner automorphism $(\alpha_f)_f \in \mathcal{Z}_{\mathbb{C}}(C)$ with $\alpha \text{id}_C = \alpha$.
- Notice that **Group** is the category of (set-based) *models* of an *algebraic theory*, i.e. a set of equational axioms between terms, namely the theory \mathbb{T}_{Grp} of groups. So **Group** = $\mathbb{T}_{\text{Grp}}\mathbf{mod}$.
- We will generalize ideas from the proof of Bergman's Theorem to give a 'syntactic' characterization of the (extended) inner automorphisms of $\mathbb{T}\mathbf{mod}$, i.e. of the covariant isotropy group of $\mathbb{T}\mathbf{mod}$, for any so-called *quasi-equational* theory \mathbb{T} .

Covariant Isotropy

- We will then use this result to characterize the covariant isotropy groups of the category **StrMonCat** of strict monoidal categories and any presheaf category **Set** ^{\mathcal{J}} .
- In particular, we will show that the covariant isotropy group of **StrMonCat** sends any strict monoidal category to its *Picard group*, i.e. its group of \otimes -invertible objects.

Quasi-Equational Theories

- What is a quasi-equational theory? (Also known as: partial Horn theory, essentially algebraic theory, cartesian theory, finite limit theory.)
- First, we need the notion of a *signature* Σ , which consists of a non-empty set Σ_{Sort} of *sorts*, and a set Σ_{Fun} of (typed) *function/operation symbols*.
- For example, the signature for *groups* has one sort X and three function symbols $\cdot : X \times X \rightarrow X$, $^{-1} : X \rightarrow X$, and $e : X$. The signature for *categories* has two sorts O, A and four function symbols **dom**, **cod** : $A \rightarrow O$, **id** : $O \rightarrow A$, and $\circ : A \times A \rightarrow A$.

Quasi-Equational Theories

- We can then form the set **Term**(Σ) of *terms* over Σ , constructed from variables and function symbols, as well as the set **Horn**(Σ) of *Horn formulas* over Σ , which are finite conjunctions of equations between terms.
- A *quasi-equational theory* over a signature Σ is then a set of *implications* (the *axioms* of \mathbb{T}) of the form $\varphi \Rightarrow \psi$, with $\varphi, \psi \in \mathbf{Horn}(\Sigma)$ (see [3]).
- The operation symbols of a quasi-equational theory are only required to be *partially* defined. If t is a term, we write $t \downarrow$ as an abbreviation for $t = t$, meaning ‘ t is defined’.

Examples

- Any *algebraic* theory, whose axioms all have the form $\top \Rightarrow \psi$, where \top is the empty conjunction. E.g. the theories of sets, semigroups, (commutative) monoids, (abelian) groups, (commutative) rings with unit, etc.
- The theories of categories and groupoids. E.g. two of the axioms of the theory of categories are

$$g \circ f \downarrow \Rightarrow \mathbf{dom}(g) = \mathbf{cod}(f),$$

$$\mathbf{dom}(g) = \mathbf{cod}(f) \Rightarrow g \circ f \downarrow.$$

- The theory of strict monoidal categories, and the theory of presheaves $\mathcal{J} \rightarrow \mathbf{Set}$ on a small category \mathcal{J} .

The Isotropy Group of a Quasi-Equational Theory

- Fix a quasi-equational theory \mathbb{T} over a signature Σ , and let $\mathbb{T}\mathbf{mod}$ be the category of (set-based) models of \mathbb{T} .
- We will now give a *logical/syntactic* characterization of the covariant isotropy group

$$\mathcal{Z}_{\mathbb{T}} : \mathbb{T}\mathbf{mod} \rightarrow \mathbf{Group}$$

of $\mathbb{T}\mathbf{mod}$.

- Using the quasi-equational syntax of \mathbb{T} , we can define a notion of *definable automorphism* for a model M of \mathbb{T} , and the definable automorphisms of any $M \in \mathbb{T}\mathbf{mod}$ form a group $\mathbf{DefInn}(M)$.

Definable Automorphisms

- Specifically, if \mathbb{T} is single-sorted, then given $M \in \mathbb{T}\mathbf{mod}$, one can form the \mathbb{T} -model $M\langle\mathbf{x}\rangle$ obtained from M by freely adjoining an indeterminate element \mathbf{x} . Elements of $M\langle\mathbf{x}\rangle$ are congruence classes $[t]$ of Σ -terms t involving \mathbf{x} and constants from M , where two terms s, t are congruent if they are provably equal in the *diagram theory* $\mathbb{T}(M, \mathbf{x})$ of M extended by the axiom $\mathbb{T} \vdash \mathbf{x} \downarrow$.

- An element $[t] \in M\langle\mathbf{x}\rangle$ is (*substitutionally*) *invertible* if there is some $[s] \in M\langle\mathbf{x}\rangle$ with

$$\mathbb{T}(M, \mathbf{x}) \vdash t[s/\mathbf{x}] = \mathbf{x} = s[t/\mathbf{x}].$$

- If f is an n -ary function symbol of Σ , then $[t] \in M\langle\mathbf{x}\rangle$ *commutes generically with f* if $\mathbb{T}(M, \mathbf{x}_1, \dots, \mathbf{x}_n)$ proves the sequent

$$f(\mathbf{x}_1, \dots, \mathbf{x}_n) \downarrow \vdash t[f(\mathbf{x}_1, \dots, \mathbf{x}_n)/\mathbf{x}] = f(t(\mathbf{x}_1), \dots, t(\mathbf{x}_n)).$$

Definable Automorphisms

- If f is an n -ary function symbol of Σ , then $[t] \in M\langle \mathbf{x} \rangle$ *reflects definedness of f* if $\mathbb{T}(M, \mathbf{x}_1, \dots, \mathbf{x}_n)$ proves the sequent

$$t[f(\mathbf{x}_1, \dots, \mathbf{x}_n)/\mathbf{x}] \downarrow \vdash f(\mathbf{x}_1, \dots, \mathbf{x}_n) \downarrow .$$

- We define **DefInn**(M) to be the group of all elements $[t] \in M\langle \mathbf{x} \rangle$ that are substitutionally invertible and commute generically with and reflect definedness of every function symbol of Σ .
- If \mathbb{T} is multi-sorted, one can extend the above definitions appropriately.

The Isotropy Group of a Quasi-Equational Theory

We then proved:

Theorem ([2])

Let \mathbb{T} be a quasi-equational theory. For any $M \in \mathbb{T}\mathbf{mod}$, the covariant isotropy group $\mathcal{Z}_{\mathbb{T}}(M)$, i.e. the group of extended inner automorphisms of M , is isomorphic to the group $\mathbf{DefInn}(M)$ of definable automorphisms of M .

Strict Monoidal Categories

- If \mathbb{T} is the quasi-equational theory of strict monoidal categories, then we proved in [2] that for any strict monoidal category \mathbb{C} , the group $\mathbf{DefInn}(\mathbb{C})$ consists (up to isomorphism) of exactly the *inner* automorphisms, i.e. the automorphisms $F : \mathbb{C} \rightarrow \mathbb{C}$ for which there is some \otimes -invertible object $c \in \mathbb{C}$ such that F is given by *conjugation* with c , i.e.

$$(a \in \mathbb{C}) \qquad F(a) = c \otimes a \otimes c^{-1}.$$

- We then deduced that

$$\mathcal{Z}_{\mathbb{T}}(\mathbb{C}) \cong \mathbf{Inv}(\mathbf{Ob}(\mathbb{C})),$$

the group of \otimes -invertible elements of the object monoid of \mathbb{C} , also known as the *Picard group* of \mathbb{C} .

Presheaf Categories

- We can also characterize the covariant isotropy group of a *presheaf category* $\mathbf{Set}^{\mathcal{J}}$ for a small category \mathcal{J} .
- Given a small category \mathcal{J} , we can define a quasi-equational theory $\mathbb{T}^{\mathcal{J}}$ whose models are functors $\mathcal{J} \rightarrow \mathbf{Set}$, i.e.

$$\mathbb{T}^{\mathcal{J}} \mathbf{mod} \cong \mathbf{Set}^{\mathcal{J}}.$$

Presheaf Categories

- If $F : \mathcal{J} \rightarrow \mathbf{Set}$ is a presheaf, we showed in [2] that $\mathbf{DefInn}(F)$ consists (up to isomorphism) of exactly the natural automorphisms $\alpha : F \xrightarrow{\sim} F$ induced by some element $\psi \in \mathbf{Aut}(\mathrm{Id}_{\mathcal{J}})$, in the sense that

$$(k \in \mathcal{J}) \quad \alpha_k = F(\psi_k) : F(k) \xrightarrow{\sim} F(k).$$




- It then follows that the covariant isotropy group $\mathcal{Z} : \mathbf{Set}^{\mathcal{J}} \rightarrow \mathbf{Group}$ is *constant* on the group $\mathbf{Aut}(\mathrm{Id}_{\mathcal{J}})$ of natural automorphisms of $\mathrm{Id}_{\mathcal{J}} : \mathcal{J} \rightarrow \mathcal{J}$.

Conclusions

- Bergman's *element-free* characterization of the inner automorphisms of groups can be used to *define* inner automorphisms in arbitrary categories.
- We have extended Bergman's *logical* characterization of the (extended) inner automorphisms of groups, i.e. of the covariant isotropy group of $\mathbf{Group} = \mathbb{T}_{\mathbf{Grp}}\mathbf{mod}$, to the covariant isotropy group of $\mathbb{T}\mathbf{mod}$ for any quasi-equational theory \mathbb{T} : we have $\mathcal{Z}_{\mathbb{T}}(M) \cong \mathbf{DefInn}(M)$ for any $M \in \mathbb{T}\mathbf{mod}$.
- Using this characterization, we have obtained logical descriptions of the definable and (extended) inner automorphisms in $\mathbf{StrMonCat}$ and presheaf categories.

Thank you!

References

-  G. Bergman. An inner automorphism is only an inner automorphism, but an inner endomorphism can be something strange. *Publicacions Matematiques* 56, 91-126, 2012.
-  P. Hofstra, J. Parker, P. Scott. Polymorphic automorphisms and the Picard group. 6th International Conference on Formal Structures for Computation and Deduction (FSCD 2021), N. Koyayashi, Ed. Dagstuhl Publications LIPIcs, Vol. 195, FSCD 2021 (to appear).
-  E. Palmgren, S.J. Vickers. Partial Horn logic and cartesian categories. *Annals of Pure and Applied Logic* 145, 314-353, 2007.